

ヨーロッパウナギの養鰻池における性と成長に関する二,三

誌名	日本水産学会誌
ISSN	00215392
巻/号	3612
掲載ページ	p. 1224-1225
発行年月	1970年12月

農林水産省 農林水産技術会議事務局筑波産学連携支援センター
Tsukuba Business-Academia Cooperation Support Center, Agriculture, Forestry and Fisheries Research Council
Secretariat



Notes on Sex and Growth of European Eels in Freshwater Eel-rearing Ponds

Syuzo EGUSA*

(Received October 30, 1970)

A huge number of elvers of *Anguilla anguilla* were imported from France to Japan in the spring of 1969 for the purpose of meeting a great shortage of elvers of *A. japonica* in the year. Since then they have been cultured, without satisfactory results, in not a few eel-farms in Shizuoka Prefecture. In regard to the cultivation of the species the following question was raised as a basic problem by experts on eels. In *A. japonica*, elvers, when cultured in freshwater ponds, develop for the most part into males¹⁾, which normally reach a marketable size of 120 to 180 mg (roughly, 42–48 cm) within two years. Natural males exceeding a length of 50 cm are not uncommon. While the males of *A. anguilla* in natural waters are reported to rarely exceed a length of 50 cm²⁾. According to EHRENBAUM³⁾, for instance, about 99% of the males which he examined were less than 40 cm long. According to KOOPS (private communication), most of the males from natural waters are less than 100 gm in weight. So provided that *A. anguilla* produces chiefly males of small size in fishponds, we cannot but feel a doubt as to the suitability of the species for cultivation. The present investigation was made to answer this question.

Material and Method

The material used consisted of 232 specimens of *A. anguilla*, ranging in body length from 32.2 to 54.3 cm, which were reared from elvers for the period April, 1969 to October, 1970 in a pond of the Marukawa Fish Farm at Yaizu City. After the body length and body weight of the fish were noted, they were dissected for sex determinations, which were made macroscopically on their gonads. Moreover, with eels measuring less than 45 cm and those measuring over 45 cm and having underdeveloped ovaries, microscopic examinations were made of sections prepared from the gonads. This re-examination revealed that the macroscopic discrimination between male and female was reliable.

Results and Discussion

The results are given in Fig. 1. Of the 232 eels, 147 (63%) were males. The percentage of males, however, varied markedly according to size. Of the 155 eels measuring

* Dept. of Fish., Fac. of Agr., Univ. of Tokyo, Bunkyo-ku, Tokyo, Japan (江草周三: 東京大学農学部)

(印刷費負担)

less than 43 cm, 138 (89%) were males, while 68 (88%) of the 77 eels measuring 43 cm or over were females. This result suggests that unlike *A. japonica*, *A. anguilla* may produce females in ponds in a relatively high percentage, though the sex-ratio of the population from which the material was taken could not be determined because of practical difficulties in investigating the size frequency distribution of the population.




Fig. 1. The influence of sex on the growth and the length-weight relationship of pond-reared *Anguilla anguilla*. The curve is a graph of the equation, $W=2.0L^3$.

From Fig. 1 it will be seen that the large majority of the males can grow to over 120 gm, though it seems that their growth, on an average, stops at a size of 40 cm or so. Further it is noticed that values of condition factor of the material fish (mean, 2.01; range, 1.39–2.55) are comparatively higher than those of pond-reared individuals of *A. japonica* of similar sizes and in the same season (mean, 1.71; range, 1.22–2.37; number of samples, 210: unpublished data).

From these findings it may be tentatively concluded that *A. anguilla* has no serious cultural handicaps in connection with sex and growth.

The author's grateful thanks are due to Mr. S. MISAKI, President of the Marukawa Fish Farm, for providing materials.

References

- 1) I. MATSUI: *J. Shimonoseki Coll. Fish.* 2(2), 1–245 (1952).
- 2) L. BERTIN: *Eels. A Biological Survey.* Cleaver-Hume Press, London, pp. 192 (1956).
- 3) E. HHRENBAUM: *Der Flußaal, Anguilla vulgaris* TURK. Handbuch der Binnenfischerei Mitteleuropas, Band II, Stuttgart (1930).