

食品酵母Candida utilisを用いて調製された短鎖ローカスト ビーンガムの凍結保護活性

誌名	日本食品工学会誌 = Japan journal of food engineering
ISSN	13457942
著者名	河原, 秀久 徳野, 悠祐 阪本, 匡章 小幡, 斉 鈴木, 雅之 槇島, 聡 裏地, 達哉
発行元	日本食品工学会
巻/号	9巻4号
掲載ページ	p. 289-296
発行年月	2008年12月

農林水産省 農林水産技術会議事務局筑波産学連携支援センター
Tsukuba Business-Academia Cooperation Support Center, Agriculture, Forestry and Fisheries Research Council
Secretariat



Cryoprotective Activity of Shortened Locust Bean Gum Prepared by Using Food Yeast *Candida utilis*

Hidehisa KAWAHARA^{1,†}, Yusuke TOKUNO¹, Masaaki SAKAMOTO¹, Hitoshi OBATA¹, Masayuki SUZUKI², Satoshi MAKISHIMA², and Tatsuya URAJI²

¹Department of Life Science and Biotechnology, Kansai University, 3-3-35 Yamate-cho, Suita-shi, Osaka 564-8680, Japan

²B Food Science Co., Ltd., 24-12 Kitahama-machi, Chita-shi 478-0046, Japan

Screening various food yeast strains revealed the growth capability of *Candida utilis* NBRC1086 on the medium of locust bean gum (LBG). The composition of the LBG medium (pH 5.0) was 0.5% (w/v) LBG, 0.3% (w/v) (NH₄)₂SO₄, 0.02% (w/v) MgSO₄ · 7H₂O, 0.01% (w/v) Yeast extract, 0.15% (w/v) NaH₂PO₄, 0.1% (w/v) KH₂PO₄. *C. utilis* has been cultivated for 24 hours, degraded by producing the β-mannosidase into the culture broth when transferred to shorten LBG (120 kDa) from LBG (300 kDa). Each LBG could be separated by gel filtration chromatography. The homogeneity of shortened LBG was confirmed by paper chromatography and gel permeation chromatography (GPC). The mannose / galactose ratio in the acid hydrolysate in shortened LBG was found to be 2.8 : 1 using HPLC. The purified shortened LBG had a high level of cryoprotective activity (100%) at a concentration of 50 μg · ml⁻¹. Furthermore, the activity of the purified shortened LBG was higher than those of LBG, mannan and other sugars. This cryoprotective activity was specified for lactate dehydrogenase (LDH) and alcohol dehydrogenase (ADH). The CP₅₀ of purified shortened LBG was 8.3 × 10 nM, which was almost one-fifth of the CP₅₀ of BSA. The cryoprotective activity of shortened LBG against a freeze-labile enzyme, had the possibility of protecting protein component in food materials, e.g. eggs, meats etc. from freezing damage, then shortened LBG could provide for various frozen foods such as good appetite feelings. The shortened LBG is expected to be more effective as a cryoprotectant than LBG for various frozen foods.

Key words: cryoprotective activity; *Candida utilis*; locust bean gum

1. Introduction

The amount of food waste for one year in Japan is 11.4 million ton in 2007, and among them approximately 10% of food waste was the expired food. In order to decrease the amount of food waste by improving the shelf life with the chilled or frozen storage techniques, key materials that have anti-freeze and anti-aging degenerative properties have been desired. One of the mannoproteins from *Pichia anomala* had a high level of cryoprotective activity [1]. Galactomannan that included in the cell membrane of *Saccharomyces cerevisiae*, has shown remarkable effects in the prevention against the frozen degeneration of food materials [1]. Galactomannans are natural nonionic heteropolysaccharide found in the endosperm of certain leguminous seeds [2]. As they can perform storage, water-retaining, and defense function in plants, they are water-

soluble and form highly viscous, stable solutions that are widely used as thickeners and stabilizers in the food, cosmetics, and textile industries [3]. Galactomannan consist of a β-1,4-linked D-mannan backbone with a D-galactopyranosyl residues α-linked to mannose C-6 sites. The degree of polymerization of the backbone is 1,000–1,500 mannose units [4]. The individual galactomannans differ from each other in their mannose/galactose residues along the main mannan chain [2].

Among some galactomannans, the most important galactomannans are locust bean gum (LBG) originated from *Ceratonia siliqua* and guar gum (GG) originated from *Cyamopsis tetragonolobus*. Mannose/galactose ratios of LBG and GG are approximately 3.5 : 1 and 1.5 : 1, respectively [5]. Bakers and Whistler reported that side-chain units of GG were disposed alternately along the mannan backbone, whereas those of LBG were disposed in uniform blocks [6]. Daas *et al.* suggested that GG has a block wise distribution of galactose side groups and LBG can show random, block wise, and ordered distributions [5].

These different structures might be caused by the highly viscosity and cryogelation in LBG. This high viscosity was resulted in difficult use at concentrations over 1% (w/v).

Though LBG is usually used as the stabilizers for ice cream, there is no data for the cryoprotective activity of LBG. If the molecular weight and configuration of commercial LBG are controlled, the anti-frozen degeneration properties must have been emphasized. However, dissolubility of galactomannan is too low to control its molecular weight and molecular configuration by using enzyme and chemical reaction. To overcome the defect of this consumption the length of the mannan chain in LBG must be shortened using β -mannanase (*endo*- β -1,4-D-mannanase EC. 3.2.1.78) and/or β -mannosidase (*exo*- β -1,4-D-mannanase EC. 3.2.1.25). β -Mannanase, which are isolated from various plants, fungi, and bacteria [7-9], is a hydrolase that catalyzes the *endo*-hydrolysis of β -1,4-mannosidic linkages in the main chain of mannans. It is extremely effective from in safety to use food yeast in a manufacturing process, however β -Mannanase originating from food yeast has briefly been little reported on.

In this paper, we described that the screening of food yeast capable of degrading LBG. We show the purification of shortened LBG and its function as a cryoprotectant having high levels of cryoprotective ability. Furthermore we suggest that shortened LBG had a low level of viscosity by comparison of the viscosity of LBG, and the suggested surface tension.

2. Materials and Methods

2.1 Microorganism and culture medium

The 21 strains of food industrial yeast belonging to the genus *Candida*, 19 strains of the genus *Saccharomyces*, 8 strains of the genus *Zygosaccharomyces*, 4 strains of the genus *Pichia*, 3 strains of the genus *Debaromyces*, 3 strains of the genus *Yarrowia*, 2 strains of the genus *Kluyveromyces*, 2 strains of the *Lipomyces* was, 2 strains of the genus *Saccharomycopsis*, 1 strain of the genus *Torulospora* and 1 strain of the genus *Yamadazyma* were purchased from NBRC culture collection (Chiba, Japan). One loopful of cells grown on a YM medium plate containing 1.5% agar, medium composition being: 1% D-glucose, 0.5% Peptone, 0.3% Yeast extract (Becton Dickinson, NJ, USA), and 0.3% Malt extract (Becton Dickinson) was inoculated into 10 ml of the YM medium (pH 5.6). The liquid culture was grown on a reciprocal shaker (120 rpm) for 1 day at 25°C, usually in Sakaguchi flasks (500 ml) containing 100 ml of the YM medium (pH 5.6).

2.2 Screening of LBG degrading food yeast and optimum culture condition

The synthetic medium (pH 5.0) containing 0.3% (w/v) $(\text{NH}_4)_2\text{SO}_4$, 0.02% (w/v) $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01% (w/v) Yeast extract (Becton Dickinson), 0.15% (w/v) NaH_2PO_4 , 0.1% (w/v) KH_2PO_4 , and 1.0% guar gum, GG (Wako Chemical Co.) or locust beam gum, LBG (Wako Chemical Co.) as carbone sources was used for the screening of LBG degrading food yeast. After 24 h cultivation at a pH of 5.0, each growth and pH were checked. In the medium containing LBG as the sole carbone source, the optimum culture conditions for the production of shortened LBG was confirmed by examining the effect of the concentration of LBG, the initial pH and the culture time.

2.3 Assay of cryoprotective activity against LDH

Cryoprotective activity was measured by a modification of Tamiya *et al.* [10, 11]. A solution of the freeze-labile enzyme, LDH (pig heart LDH obtained from Oriental Yeast Co. Ltd., Japan), was prepared in buffer A (16.66 $\text{U} \cdot \text{ml}^{-1}$). 20 μl of this LDH solution was kept in a plastic microtube and 80 μl of each sample in buffer A was added. The mixtures were then cooled under controlled conditions ($-1^\circ\text{C} \cdot \text{min}^{-1}$) and frozen at -20°C for 24 h, then thawed by controlled warming ($1^\circ\text{C} \cdot \text{min}^{-1}$). The mixture (60 μl) was then added to 3 ml of the assay mixtures (25°C), which contained 80 mM Tris-HCl buffer (pH 7.5), 100 mM KCl, 2 mM pyruvic acid, and 0.3 mM NADH. The decrease in absorbance at 340 nm was measured at 25°C using a spectrophotometer (U-2001, Hitachi Co., Japan). The residual activity of LDH after the freezing and thawing treatment was estimated by the change in absorbance during the first 2.5 min. One unit of enzyme activity was defined as the amount of enzyme catalyzing the oxidation of 1.0 μmol of NADH per min under the assay conditions. Cryoprotective activity [%] was defined by the ratio of the residual activity to decreased activity after treating with freeze and thawing. Cryoprotective activity [%] was estimated by the following equation.

$$\text{Cryoprotective activity [\%]} = \frac{\{\text{residual LDH activity [U] with sample}\} - \{\text{residual LDH activity [U] without sample}\}}{\{\text{native LDH activity [U]}\} - \{\text{residual LDH activity [U] without sample}\}} \times 100$$

2.4 Analyses of the structure and molecular weight of shortened LBG

The purified shortened LBG was separated using twice Sephacryl S-300 ($\phi 1.6 \times 60$ cm) equilibrated with 10 mM

potassium phosphate buffer (pH 7.0) containing 0.15 M NaCl. The purified shortened LBG was hydrolyzed in 2 M HCl solution at 100°C for 4 h. After hydrolyzing, this hydrolysate was neutralized with saturated BaCO₃ and then analyzed using the TOSOH sugar analysis system with a TSK gel Sugar AXG (4.6 mm ID×15 cm). Detection was accomplished by the measurement of the excitation wavelength at 288 nm and the emission wavelength at 470 nm. All sugars in the hydrolysate were converted to fluorescent using the benzimidazole and NaOH. The molecular weight of the shortened LBG was estimated with the gel filtration column, Superose 12 (ϕ 1.6×100 cm) and Sephacryl S-300 (ϕ 1.6×60 cm) using some dextrans having different molecular weight (40–190 kDa). The detections of all sugar amounts were assayed with phenol-sulfuric acid [12]. The high voltage paper chromatography to check the homogeneity of purified shortened LBG was performed by the method described by Fuller [13]. Also, the identification of shortened LBG and the estimations of the molecular weights were performed using Shodex GPC-101.

2.5 Assay of β -1,4-mannase and β -1,4-mannosidase

The β -1,4-mannanase activity in the culture broth was measured by using mannan (*S. cerevisiae*, Wako Co.) as a substrate. One unit of enzyme activity was defined as the amount of enzyme releasing 1 μ mol of reducing sugar per min [14]. The β -mannosidase activity in the culture broth was measured by using *p*-nitrophenyl- β -D-mannopyranoside (*p*Np-Man, Sigma) as a substrate. One unit of enzyme activity was defined as the amount of enzyme releasing 1 μ mol of *p*-nitrophenol per min [15]. The protein concentration was measured by the method of Bradford's method using bovine serum albumin (BSA) as the standard [16].

2.6 Assay of ice-nucleating activity

The ice nucleating activity was measured using the droplet freezing assay described by Vali [17] with a freezing nucleus spectrometer (thermoelectric plate Mitsuwa model K-1, Yamamoto Technical, Japan). Thirty drops at 10 μ l each were placed on a controlled-temperature surface and the temperature was slowly lowered from 10°C to -20°C at a rate of 1°C/min. The temperatures required to freeze 10% (T_{10}), 50% (T_{50}), and 90% (T_{90}) of the drops was measured.

2.7 Measurement of surface tension and viscosity

The surface tension of the purified shortened LBG was determined by the pendant drop method at 25°C, which was performed using an apparatus consisting of an automatic interfacial tensionmeter (CBVP-43, Kyowa Interface Science, Japan). A concentration of 1% (w/v) LBG or shortened LBG was dissolved in water. Viscosity in these solutions was performed by using the Brookfield Viscometer OV-1 at 30°C.

2.8 Chemicals

All enzymes, which were highly purified, containing lactate dehydrogenase, alcohol dehydrogenase, malate dehydrogenase, isocitrate dehydrogenase, and recombinant isocitrate dehydrogenase were purchased from the company Oriental Yeast Co. Ltd., Japan. The other chemicals were purchased from Wako chemicals Co. Ltd., Japan.

3. Results and Discussion

3.1 Screening of LBG degrading food yeast and the optimum condition for its growth

Using the screening medium containing 1.0% (w/v) GG or 1.0% (w/v) LBG as a sole carbon source, the galactomannan degrading food yeast strain was screened among the 21 strains. Although GG has a high viscosity, *Candida utilis* NBRC 1086 could grow on the GG medium after cultivation for 24 h. Next the effects of LBG concentration, pH, and cultivation time on the growth were examined. The growth of strain 1086 was made quicker by increasing the additional concentration of LBG until raised by only 1% (w/v) LBG. Optical density at 660 nm of the cell that attained its stationary phase after culturing for 20 h was 8.0. However, strain 1086 could not grow on LBG medium containing excess 1.5% (w/v) LBG causing its viscosity and solubility. Among each medium at each pH between 2 and 9, the optimum pH for its growth was 5.0. Judging from the decrease of viscosity in the culture broth and increase of reducing sugar, the optimum culture time for the growth and the production of shortened LBG was 24 hours. Some microorganisms capable of growing in a medium containing galactomannan like LBG could produce only β -mannanase [8,18], or β -mannanase and β -mannosidase [9]. After remaining in a centrifuge at 10,800×g, for 30 min to remove the cell, the resulting supernatant was examined in the presence of β -mannanase and/or β -mannosidase. Strain 1086 could produce only β -mannosidase in the culture broth, and its

activity in the culture broth was 0.52 U/mg. As β -mannosidase is an *exo*-type enzyme, the remaining LBG in the culture broth might be both whole LBG and shortened LBG.

3.2 Cryoprotective activity of LBG from the culture broth and separation of shortened LBG from the culture broth

One of the glucanase extractable mannoproteins from *Pichia anomala* had a high level of cryoprotective activity [1]. We predicted that mannan-chain had more hydrophobicity than the protein section in its mannoprotein. The resulting supernatant was fractionated using acetone. After dialyzing in distilled water using dialysis membrane (10 kDa-molecular-weight cutoff), the cryoprotective activity of the crude extract was examined. As shown in Table 1, the cryoprotective activities of the crude extract and LBG solution, which was dialyzed in distilled water, at a concentration of $100 \mu\text{g}\cdot\text{ml}^{-1}$ were 100% and 45.3%, respectively. It was suggested that LBG could hold water in the ice cream as a stabilizer [19]. Next the shortened LBG was separated using Sephacryl S-300 (Fig. 1). The enhanced activity of the shortened LBG was in a fraction indicated by the hooked symbol. This active fraction was pooled and dialyzed in distilled water, and then rechromatographed with the same condition for the isolation of purified shortened LBG. As shown in Fig. 2, the crude extract was separated into two spots that were LBG and shortened LBG. The sample pulled after separation by twice gel chromatography was only one spot. As shown in Figs. 1 and 3, its molecular weight was estimated to be approximately 120 kDa using the Sephacryl S-300 and GPC analysis. Also, the purity of separated shortened LBG could be confirmed by GPC analysis. The production of

shortened LBG having this molecular weight might to be caused by the activity of β -mannosidase.




Fig. 1 Separation of shortened LBG using gel filtration chromatography. The hooked symbol indicates the fractions of shortened LBG. The separation was performed with Sephacryl S-300 ($\phi 1.6 \times 60$ cm) with 10 mM potassium phosphate buffer (pH 7.0) containing 0.15 M NaCl.




Fig. 2 High voltage paper chromatography of purified shortened LBG.

*Purified shortened LBG was prepared with second gel filtration chromatography.

Table 1 Cryoprotective activities of various sugars.

Compound	Cryoprotective activity (%)
Sucrose	18.2
Galactose*	0
Mannose*	0
Dextran*	17.9
Mannan*	22.0
LBG*	45.3
Crude extract*	100.0
Purified S-LBG**	100.0

Sucrose concentration is 10 mg/ml.

* Sugar concentration is 100 $\mu\text{g}/\text{ml}$.

**Sugar concentration is 50 $\mu\text{g}/\text{ml}$. S-LBG indicates purified shorten LBG.

Mannan is the cell wall component from *S. cerevisiae*.




Fig. 3 GPC analysis of purified shortened LBG. Dash-dotted line is LBG, broken line is crude shortened LBG, and solid line is purified shortened LBG.

3.3 Cryoprotective activity and characterization of purified shortened LBG

Fifty $\mu\text{g}\cdot\text{ml}^{-1}$ of the purified shortened LBG had 100% (Table 1). The effect of concentration on the cryoprotective activity of shortened LBG was shown in Fig. 4. This activity sharply increased over a concentration of $5\mu\text{g}\cdot\text{ml}^{-1}$. This activity was not invariable activity over $50\mu\text{g}\cdot\text{ml}^{-1}$. The activity decreased 55% at a concentration of $100\mu\text{g}\cdot\text{ml}^{-1}$, and then increased again $200\mu\text{g}\cdot\text{ml}^{-1}$. This activity curve was the same sigmoid curve as some cryoprotective proteins [10, 20-22]. The interaction of the shortened LBG molecule with each other might differ at $50\mu\text{g}\cdot\text{ml}^{-1}$ and $100\mu\text{g}\cdot\text{ml}^{-1}$ concentrations. Following this, we examined the constituent rate of galactose and mannose in shortened LBG. Through the analysis of acid hydrolyzate using the HPLC system, we found that the ratio of mannose of the shortened LBG to the galactose of the shortened LBG was 2.8 : 1, which was different from that of the normal LBG [5]. This constituent ratio of mannose and galactose has come from the action of β -mannosidase. The cryoprotective activity of purified shortened LBG was higher than those of galactose, mannose, dextran and mannan from the cell wall of *S. cerevisiae* (Table 1). Also, the shortened LBG had no activity (0%) against malate dehydrogenase and isocitrate dehydrogenase which are well known as freeze labile enzymes, and this activity was specified for LDH and alcohol dehydrogenase. Furthermore, the CP_{50} of purified shortened LBG was 8.3×10^{-1} nM, which was almost one-fifth the CP_{50} of BSA (Table 3). This activity was slightly lower than that (CP_{50} was 2.1×10^{-1} nM) of the mannoprotein from *Pichia anomala* [1]. In general, mannan-chain has the coherent feature such as glucomannan, thereby being weak hydro-




Fig. 4 Cryoprotective activity of purified shortened LBG.

Cryoprotective activity was performed by the method described in the section on Methods and Materials. Statistical analysis was performed with a *t*-test. A *p* value less than 0.05 is considered significant.

phobicity. So we examined the surface tensions of LBG and purified shortened LBG. The surface tension of LBG was almost constant at 69 mN/m. In the shortened LBG the surface tension markedly decreased over 2 mg/ml and then this critical micelle concentration was 5 mg/ml. This result suggested that the surface tension of shortened LBG like surface active agent has an important role in expressing cryoprotective activity.

GG and LBG, which were consisted of a β -1,4-linked D-mannose backbone with a D-galactopyranosyl residues α -linked to mannose C-6 sites are widely used in the food, cosmetics, and textile industries. GG has a block wise distribution of galactose side groups and LBG can show random, blockwise, and ordered distributions [5]. This structural difference affects the actions of various enzymes, for example, β -mannanase and β -mannosidase. Even though the hydrolyzates of GG using β -mannanase were produced and characterized based on their rheological properties [23, 24], few studies have centered on the hydrolysis of LBG using various enzymes and microorganisms.

As its resulting all LBG was a shortened LBG having a molecular weight of about 120 kDa (Fig. 3), shortened LBG might be produced by the action of β -mannosidase, which was *exo*-type enzyme, in the presence of LBG. In the report of mannosidase from the genus *Candida*, *Candida albicans* could produce soluble α -1,2-mannosidase, which is involved in *N*-glycan processing in this strain [25].

In general, mannan-chain has the coherent feature such as glucomannan, and has weak hydrophobicity. The shortened LBG would have a higher hydrophobicity than natural LBG, thereby expressing high cryoprotective activity. In order to elucidate the factors in this reduction at a concentration of $100\mu\text{g}\cdot\text{ml}^{-1}$ (Fig. 4) the ice nucleation activities of the shortened LBG at each concentration were examined. The ice-nucleating temperature, T_{50} [$^{\circ}\text{C}$] of the shortened LBG at a concentration of 20, 100, and $250\mu\text{g}\cdot\text{ml}^{-1}$ was -13.9 , -17.4 and -14.2°C , respectively. Franks reported that the deciding factors in the formation of ice nuclei by materials included the following three conditions: similarity to the ice crystal lattice, the paucity of the surface charge, and the high hydrophobicity of the ice nuclei [26]. The interaction between shortened LBG and ice crystal might be related to this reduction. Furthermore, the purified shortened LBG had the 52 mN \cdot surface tension at a concentration of $5\text{mg}\cdot\text{ml}^{-1}$ (Fig. 5). A mannan-chain of the purified shortened LBG had a better interaction than that of natural LBG, thereby




Fig. 5 Effects of the concentration on surface tension.

The broken line shows distilled water. The closed circle shows the purified shortened LBG and the open circle is the natural LBG.

expressing high cryoprotective activity. Tween 80 known to be a detergent had protected the LDH during freezing and thawing [27]. Tween 80 protects LDH from denaturation when freezing and thawing by hindering its destructive interaction with the ice crystal [28]. Protection might be obtained when Tween 80 molecules compete with the protein for sites on the ice surface. The CP_{50} of purified shortened LBG was 8.3×10 nM, which was almost one-fifth of the CP_{50} of BSA (Table 2). The molar ratio of LDH and BSA was 1 : 1.7 for the CP_{50} . This value is similar to that of Tween 80 (1 : 1) [28]. Also, the cryoprotective activity of shortened LBG at $100 \mu\text{g} \cdot \text{ml}^{-1}$ under rapid cooling condition in liquid N_2 were 38%. This reduction of the activity was the same as that of Tween 80 [28]. If the protection mechanism of shortened LBG is the same as that of Tween 80, when shortened LBG is used on frozen pro-

Table 2 Comparisons of some cryoprotective matters.

	Molecular weight (kDa)	CP50(μg/ml)	CP ₅₀ (nM)	Reference
Sucrose	0.34	2.7×10^5	8.0×10^8	[20]
BSA	66	2.8×10	4.0×10^2	[20]
COR15	15	8.3×10^{-2}	5.6×10^0	[20]
COR85	350	1.5×10	4.3×10	[21]
COR26	159	1.6×10^4	1.0×10^5	[29]
HIC6	14.7	1.1×10^{-1}	7.4×10^0	[30]
CRP29	29	8.0×10^{-4}	2.7×10^{-2}	[22]
CRP62	62	2.0×10^{-1}	3.2×10^0	[31]
AS26k	26	2.0×10	7.7×10^2	[32]
COGP	83	1.7×10	2.1×10	[1]
S-LBG*	120	1.0×10	8.3×10	This study

CP_{50} indicated the concentration of each cryoprotectant required to produce a cryoprotective activity of approximately 50%.

* S-LBG indicates purified shorten LBG.

Table 3 Comparison of LBG and shorten LBG.

	LBG	Shorten LBG
Molecular weight (kDa)	300	120
CP_{50} (nM)	6.7×10^2	8.3×10
Viscosity (cP)	100.6	33.2

cessed foods, the frozen food might have a subsequently high level of moisture and quality. The viscosity of shortened LBG at a concentration of $1 \text{ mg} \cdot \text{ml}^{-1}$ was lower than that of natural LBG. By the reduction of viscosity, the solubility of shortened LBG was higher than that of natural LBG (Table 3).

We could find that the shortened LBG produced by *Candida utilis* NBRC 1086 had a high level of cryoprotective activity, and surface tension. Since this shortened LBG has the possibility of holding water, the use of shortened LBG might provide a novel texture for the cryogel in the mixture of other polysaccharide and shortened LBG in future.

Acknowledgment

We would like to the The Iijima Memorial Foundation For The Promotion Of Food Science And Technology for their generous financial support of our research in 2007.

References

- [1] H. Kawahara, N. Omori, H. Obata; Cryoprotective activity of mannoprotein from the cell membrane of *Pichia anomala*. *Cryo. Letters*, **29**, 437-445 (2008).
- [2] P. M. Dey; Biochemistry of plant galactomannans. *Adv. Carbohydr. Chem. Biochem.*, **35**, 341-376 (1978).
- [3] H. Maier, M. Anderson, C. Kari, K. Magnuson, R. L. Whistler. In *Industrial Gums, Polysaccharides and Their Derivatives*; R. L. Whistler, J. N. BeMiller, Eds.; Academic Press: San Diego, CA, 1993 pp.181-226.
- [4] I. C. M. Dea, E. R. Morrison: Chemistry and interactions of seed galactomannans. *Adv. Carbohydr. Chem. Biochem.*, **31**, 241-312 (1975).
- [5] P. J. H. Daas, H. Schols, H. H. J. de Jough: *Carbohydr. Res.*, **329**, 609-619 (2000).
- [6] C. W. Baker, R. L. Whistler: *Carbohydr. Res.*, **45**, 237-243 (1975).
- [7] M. Kurakake, T. Komaki: Production of β -mannanase and β -mannosidase from *Aspergillus awamori* K4 and their properties. *Curr. Microbiol.*, **42**, 377-380 (2001).
- [8] A. Sunna, M.D. Gibbs, C.W. Chin, P.J. Nelson, P.L. Bergquist: A gene encoding a novel multidomain β -1,4-

- mannanase from *Caldobacillus cellulovorans* and action of the recombinant enzyme on kraft pulp. *Appl. Environ. Microbiol.*, **66**, 664-670 (2000).
- [9] B. V. McCleary: β -D-Mannanases. *Methods Enzymol.*, **160**, 596-610 (1988).
- [10] H. Obata, N. Muryoi, H. Kawahara, A. Nishiyama: Purification and characterization of uridine phosphorylase from the ice-nucleating bacterium, *Pantoea agglomerans* NBRC12686. *Cryo.Letters*, **25**, 195-204 (2004).
- [11] T. Tamiya, N. Okahashi, R. Sakuma, T. Aoyama, T. Akahane, J. Matsumoto: Freeze denaturation of enzymes and its prevention with additives. *Cryobiology*, **22**, 446-456 (1985).
- [12] T. Masuko, A. Minami, N. Iwasaki, T. Majima, S. Nishimura, Y. C. Lee: Carbohydrate analysis by a phenol-sulfuric acid method in microplate format. *Anal. Biochem.*, **339**, 69-72 (2005).
- [13] K. W. Fuller, D. H. Northcote: A micro method for the separation and determination of polysaccharides by zone electrophoresis. *Biochem. J.*, **64**, 657-663 (1956).
- [14] B. V. McCleary: A simple assay for β -D-mannanase. *Carbohydr. Res.*, **67**, 213-221 (1978).
- [15] M. Arai, H. Fujimoto, T. Ooi, S. Ogura, S. Murao: Purification and properties of a β -mannosidase from *Aspergillus aculeatus*. *Oyo Toshitsu Kagaku*, **42**, 49-51 (1995).
- [16] M. Bradford: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.*, **72**, 248-254 (1976).
- [17] G. Vail : Quantitative evaluation of experimental results on the heterogeneous freezing nucleation of supercooled liquids. *J. Atmos. Sci.*, **28**, 402-409 (1971).
- [18] N. Takeda, K. Hirose, K. Uchimura, Y. Nogi, Y. Harada, M. Akita, R. Usami, Y. Yoshida, W. D. Grant, S. Ito, K. Horikoshi: Alkaline mannanase from a novel species of alkaliphilic *Bacillus*. *J. Appl. Glycosci.*, **51**, 229-236 (2004).
- [19] A. Regand, H. D. Goff: Effect of biopolymers on structure and ice recrystallization in dynamically frozen ice cream model systems. *J. Dairy Sci.*, **85**, 2722-2732 (2002).
- [20] C. Lin, M. F. Thomashow: A cold-regulated *Arabidopsis* gene encodes a polypeptide having potent cryoprotective activity. *Biochem. Biophys. Res. Commun.*, **183**, 1103-1108 (1992).
- [21] M. F. Thomashow: Molecular genetics of cold acclimation in higher plant. *Adv. Genet.*, **28**, 99-131 (1990).
- [22] N. Koda, T. Asaeda, K. Yamade, H. Kawahara, H. Obata: A novel cryoprotective protein (CRP) with high activity from the ice-nucleating bacterium, *Pantoea agglomerans* IFO12686. *Biosci. Biotechnol. Biochem.*, **65**, 888-894 (2001).
- [23] S. Mahammad, D. A. Comfort, R. M. Kelly, S. A. Khan: Rheological properties of guar galactomannan solutions during hydrolysis with galactomannanase and α -galactosidase enzyme mixtures. *Biomacromolecules*, **8**, 949-956 (2007).
- [24] M. Kuratake, T. Sumida, D. Masuda, S. Oonishi, T. Komaki: Production of galacto-manno- oligosaccharides from guar gum by β -mannanase from *Penicillium oxalicum* SO. *J. Agric. Food Chem.*, **54**, 7885-7889 (2006).
- [25] H. M. Mora-Montes, E. López-Romero, S. Zinker, P. Ponce-Noyola, A. Flore-Carreón : Purification of soluble α -1,2-mannosidase from *Candida albicans* CAI-4. *FEMS Microbiol. Lett.*, **256**, 50-56 (2006).
- [26] F. Franks in "Biophysics and Biochemistry at Low temperatures" Japan UNI Agency Inc. 1985, pp. 36-37.
- [27] J. F. Carpenter, M. J. Pikal, B. S. Chang, T. W. Randolph : Rational design of stable lyophilized protein formulations : some practical advice. *Pharm. Sci.*, **85**, 1325-1330 (1997).
- [28] A. Hillgren, J. Lindgren, M. Aldén: Protection mechanism of Tween 80 during freeze-thawing of a model protein, LDH. *Int. J. Pharm.*, **237**, 57-59 (2002).
- [29] H. Obata, H. Ishigaki, H. Kawahara, K. Yamade : Purification and characterization of a novel cold-regulated protein from an ice-nucleating bacterium, *Pseudomonas fluorescens* KUIN-1. *Biosci. Biotechnol. Biochem.*, **62**, 2091-2097 (1998).
- [30] K. Honjoh, H. Matsumoto, H. Shimizu, K. Oomiya, K. Tanaka, Y. Oda, R. Tanaka, T. Joh, K. Suga, T. Miyamoto, M. Ito, S. Hirano : Cryoprotective activities of group 3 late embryogenesis abundant proteins from *Chlorella vulgaris* C-27. *Biosci. Biotechnol. Biochem.*, **64**, 1656-1663 (2000).
- [31] L. G. Neven, D. W. Haskell, A. Hong, Q-B. Li, C. L. Guy : Characterization of a spinach gene responsive to low temperature and water stress. *Plant Mol. Biol.*, **21**, 291-305 (1993).
- [32] M. Momma, K. Haraguchi, M. Saito, K. Chikuni, K. Harada : Purification and characterization of the acid soluble 26-kDa polypeptide from soybean seeds. *Biosci. Biotechnol. Biochem.*, **61**, 1268-1289 (1997).

◇◇◇◇ 和文要約 ◇◇◇◇

食品酵母 *Candida utilis* を用いて調製された 短鎖ローカストビーンガムの凍結保護活性

河原秀久^{1,†}, 徳野悠祐¹, 阪本匡章¹, 小幡 齊¹, 鈴木雅之², 槇島 聡², 裏地達哉²

¹ 関西大学化学生命工学部生命・生物工学科, ² 物産フードサイエンス株式会社

種々の食品酵母株のスクリーニングテストによって、*Candida utilis* NBRC1086 がローカストビーンガム (LBG) 培地で生育できることが明らかになった。 *C. utilis* は、0.5% (w/v) LBG, 0.3% (w/v) (NH₄)₂SO₄, 0.02% (w/v) MgSO₄·7H₂O, 0.01% (w/v) Yeast extract, 0.15% (w/v) NaH₂PO₄, 0.1% (w/v) KH₂PO₄ を含む LBG 培地 (pH 5.0) で、24 時間培養された。また、*C. utilis* は、培地中に β-マンノシダーゼを生産することによって LBG (300 kDa) を短鎖 LBG (120 kDa) に分解していた。各々の LBG は、ゲルろ過クロマトグラフィによって分離された。短鎖 LBG の単一性はペーパークロマト

グラフィと GPC 分析によって確認された。短鎖 LBG の酸加水分解物のマンノースとガラクトース比は、2.8:1 であった。精製した短鎖 LBG は、50 μg/ml 濃度で高い凍結保護活性 (100%) を示し、LBG やマンナンおよび他の糖よりも高い活性であった。この凍結保護活性は乳酸脱水素酵素とアルコール脱水素酵素に特異的であった。精製短鎖 LBG の CP50 は、8.3×10 nM であり、この値は牛血清アルブミンの約5分の1であった。また、精製した短鎖 LBG は、冷凍液卵を安定化させることも確認できた。我々は、短鎖 LBG が多様な冷凍食品に対して凍結保護剤の可能性をもっていると期待している。

(受付 2008 年 7 月 31 日, 受理 2008 年 10 月 30 日)

1 〒 564-0073 大阪府吹田市山手町 3-3-35

2 〒 478-0046 愛知県知多市北浜町 24-12

† Fax: 06-6368-0832, E-mail: kawahara@ipcku.kansai-u.ac.jp